arXiv:1210.1577v1 [nucl-th] 4 Oct 2012

# Baryon Annihilation in Heavy Ion Collisions

Yinghua Pan
*Department of Physics, Harbin Institute of Technology,*
*Harbin, Heilongjiang 150006, People's Republic of China and*
*Department of Physics and Astronomy and National Superconducting Cyclotron Laboratory, Michigan State University*

Scott Pratt
*Department of Physics and Astronomy and National Superconducting Cyclotron Laboratory, Michigan State University*

(Dated: October 8, 2012)

The role of baryon-antibaryon annihilation during the hadronic stage of a relativistic heavy ion collision is explored by simulating the chemical evolution of a hadron gas. Beginning with a chemically equilibrated gas at an initial temperature of 165 MeV, it is assumed that the gas expands according to the equations of a one-dimensional Bjorken expansion. The evolution includes both annihilation and regeneration of baryons. From an initial time of 5 fm/$c$ until a final time of 20 fm/$c$, the number of baryons drops by approximately half for the case where there is no net baryonic charge. This is consistent with measurements from the ALICE Collaboration at the LHC, where the $p/\pi$ ratio is roughly 50% lower than what comes from hydrodynamic models that ignore baryon annihilation. When the calculations are performed without the inverse process, e.g. $N\pi \rightarrow p\bar{p}$, the loss of baryons was found to be significantly larger. Baryon annihilation is shown to alter the extracted chemical breakup temperature. Assuming that annihilation cross sections are independent of the strangeness and isospin of the annihilating baryon and anti-baryon, the loss of strange baryons from annihilation is found to be similar.

## INTRODUCTION

High energy heavy ion collisions provide a unique opportunity to investigate the properties of nuclear matter with high energy density, a few GeV/fm$^3$, which is several times the energy density of a proton. At these temperatures, $\gtrsim 170$ MeV, matter undergoes a transition between the hadronic phase and the quark-gluon plasma phase (QGP)[1, 2]. In the prevailing view, the violently interacting produced particles validate the use of thermodynamics [3–10] to describe the chemistry and hydrodynamics [11] to describe the evolution. A typical assumption is that the hadronic chemistry is described by chemical freeze-out at a single temperature. This temperature is also associate with the QGP-hadron transition, or hadronization. Although hadrons certainly collide a few more times after hadronization, many models typically assume such effects to be negligible. A variety of particle ratios are reasonably well fit by a single temperature and a few chemical potentials to account for non-zero conserved charges such as strangeness, baryon number and electric charge. Additionally, fugacities have been added to account for the inability of the system to produce the requisite number of up, down and strange quarks for equilibrium [12]. These pictures are all based on the idea that, due to the strong interactions between quarks, all partitions into various hadron species are thermally sampled during hadronization, and that the evolution of chemical abundances after hadronization comes only from decays.

Of course, the chemical make-up does not completely freeze-out — hadrons indeed undergo chemical reactions during the hadronic stage. The effects of such evolution have been studied in two contexts. First, microscopic transport models such as URQMD [13] include inelastic processes. Although these models include annihilation processes, e.g., $p\bar{p} \rightarrow n\pi$ [14, 15], no microscopic model has yet included the inverse process. A second class of approaches involves solving for the chemical rates of a chemically equilibrated gas. Given that there are hundreds of resonances, such models typically assign fugacities to a few numbers such as effective pion number, or the net number baryons plus antibaryons, and then solve for the evolution of the fugacities [16–19]. In Reference[19] it was shown that reformation rates, e.g. $n\pi \rightarrow p\bar{p}$, were important.

The shape of pion, kaon and proton spectra can be well described by hydrodynamic models that assume chemical equilibrium at the hadronization temperature. However, the normalization of the proton spectra tends to significantly vary from experimental measurements from PHENIX at RHIC, which tend to fall $\sim 40\%$ below model predictions that fit the yields of pions and kaons [20] when comparing to PHENIX spectra from RHIC [21]. Similar shortcomings are seen at ALICE when comparing to thermal yields [14]. Using a microscopic model, a reduction of roughly this size was seen [15], but those calculations ignore the inverse rates, which according to [19] largely cancel the annihilation rates. The goal of this paper is to solve for the chemical rates of a wide range of hadronic species where baryon annihilation and reformation are consistently included. This is similar in spirit to what was done in [19], but in more detail with a larger number of resonances considered. The goal will be to see whether in a consistent calculation the effect of annihilations might account for a $\sim 40\%$ reduction in the $p/\pi$ ratio, and to discern whether that reduction differs for hyperons.

In order to concentrate on the chemistry, we employ a very simplified model of the space-time evolution. We consider a one-dimensional boost-invariant expansion [22]. In such a model, all fluid elements expand identically and their volumes increase inversely proportional to the time as measured from the beginning of the collision. We assume that the hadronization temperature, 165 MeV [23], is reached at a time of 5 fm/$c$, and that reactions cease at 20 fm/$c$ to be roughly consistent with expectations for 200$A$ GeV Au+Au collisions. In a more realistic model, the hadronization temperature would be reached at different times depending on the transverse position, and transverse expansion would accelerate the cooling at later times. The decoupling would be gradual, and the various species would lose thermal contact with one another and not maintain local kinetic equilibrium. Nonetheless, this picture is adequate to gauge the significance of baryon annihilation.

## DESCRIPTION OF CALCULATION

We model the hadronic stage assuming that all resonances are initially populated according to chemical equilibrium according to a temperature $T_0$, set by default to 165 MeV. All chemical potentials are set to zero. The list of 319 resonances extends to masses up to 2.25 GeV. Although one should have a non-zero baryonic chemical potential at high RHIC and LHC energies, it will be ignored for this study given that the our goal is to roughly gauge the importance of annihilation effects.

After the initial time, we assume that collisions are sufficiently frequent to preserve local kinetic equilibrium, even though inelastic collisions are insufficient for maintaining chemical equilibrium. Each species is assigned a chemical potential, $\mu_i$. Each $\mu_i$ is initially set to zero, and if chemical equilibrium were maintained each $\mu_i$ would remain zero.

One can derive the density and entropy contributions for each species from a grand canonical ensemble,

$$\begin{aligned}\ln Z(T,V,\{\mu_i\}) &= \sum_{species\ i} \frac{g_i V}{(2\pi)^3}\int d^3p \ln(1 \pm e^{\frac{-(E_i-\mu_i)}{T}})^{\pm 1} \\ &= \sum_i \frac{V g_i}{2\pi^2}\sum_k (\pm 1)^{k+1}\left(\frac{m_i^2 T}{k}K_0(km_i/T) + \frac{2m_iT^2}{k^2}K_1(km_i/T)\right)e^{k\mu_i/T}.\end{aligned} \tag{1}$$

where $g_i$ and $\mu_i$ are the degeneracy and chemical potential of hadron species $i$, $E_i = \sqrt{p^2+m_i^2}$ where $m_i$ is the particle mass, and V is the hadron gas volume. Once the partition function is known, the number densities, pressure, energy density and entropy can be found by differentiation:

$$\begin{aligned} n_i &= \frac{g_i}{2\pi^2}\sum_k (\pm 1)^{k+1}\left(m_i^2 T K_0(km_i/T) + \frac{2m_iT^2}{k}K_1(km_i/T)\right)e^{k\mu_i/T}, \\ P &= \frac{T}{V}\ln Z, \\ \epsilon &= \frac{T^2}{V}\frac{\partial \ln Z}{\partial T} = \sum_i \frac{g_i}{2\pi^2}\sum_k (\mp 1)^{k+1}\left[\frac{3m_i^2T^2}{k}K_0(km_i/T) + \left(m_i^3 T + \frac{6m_iT^3}{k^2}K_1(km_i/T)\right)\right]e^{k\mu_i/T}, \\ s &= \frac{P+\epsilon-\sum_i \mu_i n_i}{T}, \end{aligned} \tag{2}$$

where $K_n$ is the Bessel function of order $n$. Bose/Fermi effects enter through considering the sum over $k$. Bose effects are neglected for this study by considering only the $k = 1$ term. By including the $k > 1$ terms, the density of pions would rise at the $\sim 5\%$ level, while the effect on other species would be negligible.

In order to maintain chemical equilibrium as the system cools, the population of heavier particles needs to decrease due to the fact that the thermodynamic penalty for mass is more important at low temperature. Even pions, the lightest mesons, need to reduce their population through annihilation and absorption to maintain chemical equilibrium [16, 24]. Since baryons are heavy, their population is most sensitive to the temperature and their populations stray furthest from equilibrium during the rapid cooling of the hadronic stage.

Since we are interested in simply gauging the effects of annihilation, we consider a simplified picture of the expansion, where densities follow a one-dimensional boost-invariant expansion [22]. This picture neglects transverse expansion and assumes that longitudinal acceleration is zero. Based on the assumption above, the continuity equations can be easily be solved in Bjorken coordinates should all chemical rates be turned off,

$$\begin{aligned} n_i(\tau) &= n_i(\tau_0)\frac{\tau_0}{\tau}, \\ s(\tau) &= s(\tau_0)\frac{\tau_0}{\tau}. \end{aligned} \tag{3}$$

Here, $\tau_0$ is the proper time in the local frame of the fluid cell when the system starts to expand and is set at $\tau_0$= 5 fm/c. The evolution equations for the chemistry are modified in the presence of chemical rates,

$$\frac{d}{d\tau}\left[\tau n_i(\tau)\right] = \tau R_i(\tau), \tag{4}$$

where $R_i$ is net production/annihilation rate per unit $d^4x$ of particles of species $i$. The net rates, $R_i$, are all zero at equilibrium, but as the system cools the system loses equilibrium and the rates become non-zero (usually negative) to push toward equilibrium.

Once the system loses chemical equilibrium, entropy can be generated,

$$\frac{d}{d\tau}\left[\tau s(\tau)\right] = -\tau \sum_i R_i(\tau)\frac{\mu_i(\tau)}{T(\tau)}. \tag{5}$$

Given the entropy and number densities, one can solve for the temperature and chemical potentials. Thus, Given the rates $R_i$, Eq.s (4-5) allow one to solve for the evolution of the number and entropy densities, or equivalently the chemical potentials and temperatures, as a function of $\tau$. A multidimensional Newton's method is used to numerically determine $\mu_i$ and $T$ from $n_i$ and $s$.

One contribution to $R_i$ is resonances decays. For the reaction $A \to a_1 + a_2 + \cdots + a_n$, the contribution to the growth rate for resonance of type $A$ is

$$R_A(\tau) = -\left(1 - e^{\frac{\mu_{a_1}+\mu_{a_2}+\cdots+\mu_{a_n}-\mu_A}{T}}\right) \langle \Gamma_{A\to a_1\cdots a_n}\rangle \, n_A(\tau). \tag{6}$$

Here $\langle\Gamma\rangle$ is the decay rate of the resonance, which is the nominal width divided by the average Lorentz time dilation factor,

$$\langle\Gamma\rangle = \frac{\Gamma}{(2\pi)^3 n_A}\int (d^3p/E_A)\; m_A f(p). \tag{7}$$

Partial widths are taken from the particle data book [25]. The contribution from the inverse reaction, $a_1 + \cdots a_n \to A$, is accounted for by the factor $\exp[(-\mu_A + \mu_1 \cdots + \mu_n)/T]$. This forces $R_A$ to go to zero when the chemical potentials balance. This channel also contributes to the growth rates for $a_1 \cdots a_n$ with the same magnitude but opposite sign. All the known hadronic decay channels from the particle data book are included in the calculation.

The second type of chemical reaction to be considered is that from baryon annihilation, $A + \bar{B} \to a_1 + \cdots a_n$. The contribution to the growth rate of $A$ in this case is

$$R_A(\tau) = -\left(1 - e^{(\mu_{a_1}+\mu_{a_2}+\cdots+\mu_{a_n}-\mu_A-\mu_B)/T}\right) \langle \sigma_{A+\bar{B}\to a_1\cdots a_n} v_{\rm rel}\rangle \, n_A(\tau) n_B(\tau). \tag{8}$$

The average $\langle\sigma v_{\rm rel}\rangle$ for a relativistic gas is

$$\langle\sigma v_{\rm rel}\rangle = \frac{1}{(2\pi)^6 n_A n_B}\int (d^3p_a/E_a)(d^3p_b/E_b)\; f_a(p_a) f_b(p_b)\sigma(s)\sqrt{(p_a\cdot p_b)^2 - m_a^2 m_b^2}. \tag{9}$$

We apply a simplified form for the $p\bar{p}$ annihilation cross section that is accurate for $P_{\rm lab} > 100$ MeV/$c$ [26, 27],

$$\sigma = 67\; P_{\rm lab}^{-0.7}{\rm mb}. \tag{10}$$

Without particularly good justification, we further assume all baryon-antibaryon pairs have the same annihilation cross section as $p\bar{p}$. Averaged over momentum, the average $\langle\sigma v_{\rm rel}\rangle$ is near 5 fm·$c$ for $p\bar{p}$. Other inelastic processes, e.g., $\Delta + N \to NN$, are ignored for this study as we are focusing on baryons. Such processes can change $n_\pi$ at the level on the order of 10% [16, 28].

Experimental data show that the most likely number of pions in $p\bar{p} \to N\pi$ is $\sim 5$, and to simplify the calculation we assume that all annihilations proceed through the 5 $\pi$ production channel. For pair annihilation with nonzero strangeness, we employ the minimal rule for kaon production, i.e, the number of kaons in the final state equals the initial strangeness. For example, if the initial strangeness is unity, then final production of annihilation will have one kaon. We still assume final production has five particles on average. The number of pions, $n_\pi$, will be $5 - n_K$.

Before displaying numerical results, we summarize the assumptions in our calculation. We consider a system with initial charge set to zero, and at chemical equilibrium so that $\mu_i = 0$ for all species. The initial temperature is set to 165 MeV, and the initial time is set to $\tau_0 = 5$ fm/$c$. The calculation proceeds until freezeout at 20 fm/$c$. Only decays through the strong interaction are considered since both electromagnetic and weak decays are negligible on these time scales. The calculation considered 319 resonances, with masses up to 2.25 GeV/$c^2$.

## RESULTS

Figure 1 shows the evolution of the densities of pions, kaons, protons, Lambdas, Sigma, Cascade and Omega baryons. Results are displayed both for when annihilations are enabled and disabled. When annihilations are disabled, the populations of stable particles increase due to feeding from the decays of unstable particles. Decays increase the number of protons by roughly 50%, mainly through delta decays. The effect of baryon annihilation is to lower the baryon yields by ∼40%, to increase pion populations by ∼10% and to increase the kaon yields by ∼5%. Annihilations reduce the yields of strange baryons by approximately the same factor as protons. For the times immediately after $\tau_0$, annihilations play no role because the inverse process exactly cancels annihilation at equilibrium. However, within a few fm/$c$, the system is no longer at chemical equilibrium and the curves with and without annihilation diverge.

As the system cools, chemical equilibrium is lost and the chemical potentials become non-zero. Figure 2 shows the evolution of the temperature and chemical potentials as a function of time. The temperatures are slightly enhanced



FIG. 1. Hadronic densitiies, scaled by $\tau$, for the $\pi$ and $K$ mesons, and for the $p$, $\Lambda$, $\Sigma$, $\Xi$, and $\Omega$ baryons. In the absence of annihilation, the proton yield increases due to the decay of resonances like the $\Delta$. By adding annihilation, all baryon populations fall by $\sim 40\%$. Meson populations are modestly increased by the annihilation processes.

by annihilation, because the process $p\bar{p} \rightarrow 5\pi$ creates pions with higher average energy than the thermal value. The chemical potentials develop more strongly for more massive particles since their populations are more sensitive to temperature. The lower panel demonstrates the degree to which the inverse processes cancel annihilation. At the initial time, chemical populations are equilibrated and the inverse processes, e.g. $5\pi \rightarrow p\bar{p}$ exactly cancel the annihilation processes. By the end of the reaction, the inverse processes cancel $\sim 55\%$ of the annihilations. This underscores the importance of considering the inverse processes.

A common use of particle ratios is to determine the chemical freeze-out temperature. Although analyses of this type involve a global fit, the temperature is mainly determined by ratios between the heaviest and lightest particles



FIG. 2. The evolution of the temperature and chemical potentials for three species as a function of time in the upper two panels. The lower panel shows the fraction of the annihilation rate that is uncanceled by the inverse process. Initially, the system is at thermal equilibrium and annihilation is perfectly canceled by the inverse process, e.g., $5\pi \rightarrow p\bar{p}$. By the time breakup occurs, the inverse process cancels ~ 55% of the annihilations.

since they are the most sensitive to the temperature. The $p/\pi$ ratio is especially important. Since such analyses, c.g. [6], ignore baryon annihilation it is instructive to see the degree to which annihilation processes distort the extraction of a chemical freezeout temperature. Figure 3 displays the final $p/\pi^+$ ratio as a function of the initial temperature for calculations with and without baryon annihilation. The ALICE collaboration at the LHC[29] measured $p/\pi^+ = 0.46 \pm 10\%$, and is illustrated with a grey band in Fig. 3. Whereas that would correspond to a chemical freeze-out temperature of 145 MeV if baryon annihilation were ignored, one would infer a breakup temperature near 170 MeV if annihilation were considered.

## CONCLUSIONS

Figure 1 shows that annihilation processes during the hadronic stage play a significant role in the baryo-chemistry of the quark-gluon plasma. Calculated baryon yields that include the effects of annihilation are reduced by ~ 40%. This helps explain the magnitude of the $p$ and $\bar{p}$ spectra at the LHC relative to the pion spectra without surrendering



FIG. 3. The final $p/\pi^+$ ratio as a function of $T_0$, the initial temperature of chemical freezeout. Calculations with and without baryon annihilation differ substantially. If one were to use the experimental $p/\pi^+$ ratio measured by ALICE to infer $T_0$, including annihilation would change the inferred value from $\sim 145$ MeV to $\sim 173$ MeV.

the assumption of chemical equilibrium at the advent of the hadronic stage. Figure 1 also displays results from calculations where the inverse reactions, e.g. $5\pi \rightarrow p\bar{p}$, were not included. In this case baryon yields would have fallen by another 40%. These results show both the importance of including annihilations, and when doing so, to consistently include the inverse reactions.

The population of strange baryons is similarly affected. There are significant uncertainties in the experimental measurement of strange baryons, and it is premature to state whether the yields are suppressed by amounts similar to what is observed for protons. If it turns out that hyperon yields are not similarly suppressed, it may inspire one to consider whether annihilation cross sections involving strange baryons differ from those for non-strange baryons.

Although sufficient for obtaining an understanding of the importance of annihilation chemistry during the hadronic stage, the model employed here is too simplistic to compare to data at even the 10% level. The assumptions are numerous: hadronic equilibrium at the onset of hadronization, one-dimensional boost-invariant dynamics, and local kinetic equilibrium that abrubtly ends at $\tau = 20$ fm$/c$. Further, some of the chemistry is neglected. Inelastic collisions, such as $\Delta N \rightarrow NN$ can lower the final meson populations on the order of 10%. To improve the model to the 10% level, one should employ a microscopic simulation of the hadronic stage. This can account for the three-dimensional expansion, provide a realistic freeze-out picture, and allow for the loss of local kinetic equilibrium. However, the current generation of such models neglects reformation processes, as well as some of the inelastic processes that affect the final chemical make-up measured in experiment. For instance, although baryon annihilation processes can be enabled in URQMD [13], the inverse processes are not. Reactions where $n > 2$ particles collide are more difficult to simulate in microscopic models. If one wishes to increase the reliabilities of these models to the ten percent level, such improvements are necessary.

Figure 1 also showed the effect of annihilation on strange baryon populations. However, these results are not trustworthy given the lack of knowledge of annihilation cross sections for hyperons. Whereas the nucleon-antinucleon cross sections does not strongly depend on isospin, e.g. $\bar{p}p$ and $\bar{p}n$ annihilation cross sections are similar [30], little is known about the annihilation cross sections for hyperons, either with antinucleons or antihyperons, and one must resort to unconstrained model calculations [31] or to fits to particle yields in heavy ion collisions [26]. Without direct experimental information about antihyperon annihilation cross sections, interpreting the yields of strange baryons will be problematic. In a future study, hyperon annihilation will be studied in more detail. First, it will investigated whether changes in hyperon annihilation cross sections significantly affect the hyperon survival probability. Secondly, the process where kaons scatter inelastically off nucleons to form hyperons will be investigated. In our calculations we see that $\mu_K + \mu_p > \mu_\Lambda + \mu_\pi$. Thus, inelastic 2 → 2 scatterings have the potential to lower nucleon yields while enhancing hyperon yields. In fact, such a process would successively increase the yields of hyperons of increasing strangeness. Although the trend is clear, the magnitude of the effect is not yet understood.

[1] G. Van Buren for the STAR Collaboration, in Proceedings of Quark Matter 2002, Nantes, France, 2002,edited by H. Gutbrod, J. Aichelin, and K. Werner (Elservier, Amsterdam,2003).
[2] L. McLarren, Rev. Mod. Phys. **58**,1001 (1986).
[3] J. Stachel and P. Braun-Munzinger, Phys. Lett. B **216**, 1 (1989).
[4] P. Braun-Munzinger, J. Stachel, J. Wessels, and N. Xu, Phys. Lett. B **344**, 43 (1995).
[5] P. Braun-Munzinger, J. Stachel, J. Wessels, and N. Xu, Phys. Lett. B **365**, 1 (1996).
[6] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, J. Phys. G G **38**, 124081 (2011) [arXiv:1106.6321 [nucl-th]].
[7] J. Rafelski, Phys. Lett. B **262**, 333 (1991).
[8] N. J. Davidson, H. G. Miller, R. M. Quick, and J. Cleymans, Phys. Lett. B **255**, 105 (1991).
[9] J. Sollfrank, M. Gazdzicki, U. Heinz, and J. Rafelski, Z. Phys. C **61**, 659 (1994); U. Heinz, Nucl. Phys. A **566**,205c (1994).
[10] S. Wheaton and J. Cleymans, hep-ph/0407174.
[11] U. heinz, arXiv: 0901.4355v1 [nucl-th] (2009).
[12] G. Torrieri, S. Jeon and J. Rafelski, Phys. Rev. C **74**, 024901 (2006) [nucl-th/0503026].
[13] S.A.Bass et al [URQMD], Progr. Part. Nucl. Physics Vol. **41**, 225 (1998).
[14] J. Steinheimer, V. Koch and M. Bleicher, arXiv:1207.2791 [nucl-th].
[15] I. .A. Karpenko, Y. .M. Sinyukov and K. Werner, arXiv:1204.5351 [nucl-th].
[16] S. Pratt, K. Haglin, Phys. Rev. C **59**, 3304 (1999).
[17] Chung-sik Song and V. Koch, Phys. Rev. C **55**, 3026 (1997).
[18] R. Rapp, Phys. Rev. C **66**, 017901 (2002).
[19] R. Rapp and E. V. Shuryak, Phys. Rev. Lett. **86**, 2980 (2001) [hep-ph/0008326].
[20] S. Pratt, Phys. Rev. C **85**, 014904 (2012).
[21] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. C **69**, 034909 (2004) [nucl-ex/0307022].
[22] J. D. Bjorken, Phys. Rev. D **27**,140151 (1983).
[23] L. Kumar, arXiv:1109.5313v1 [nucl-ex].
[24] C. Greiner, C. Gong and B. Muller, Phys. Lett. B **316**, 226 (1993) [hep-ph/9307336].
[25] Particle Data Group, *The Review of Particle Physics*, http://pdg.lbl.gov (2012).
[26] G. J. Wang, R. Bellwied, C. Pruneau, G. Welke, arXiv:nucl-th/9806006v1.
[27] A. Bianconi et al., Phys. Lett. B481, 194 (2000).
[28] Bao-An Li, Che Ming Ko, Phys. Rev. C **52**, 2037 (1995).
[29] B. Abelev *et al.* [ALICE Collaboration], arXiv:1208.1974 [hep-ex].
[30] T. Elioff, L. Agnew, O. Chamberlain, H. M. Steiner, C. Wiegand, T. Ypsilantis, Lawrence Berkeley National Lab. Publication, http://escholarship.org/uc/item/1046p96v (1961).
[31] J.S. Kovacs and D.B. Lichtenberg, Nuovo Cimento **10**, 376 (1959).